



Navigating prior authorizations and appeals for DUPIXENT® (dupilumab)

**For patients with moderate-to-severe eosinophilic
or oral corticosteroid-dependent asthma**

- ⋮ This guide provides information about health plan requirements that may be
- ⋮ required when submitting prior authorizations (PAs) or appealing PA denials
- ⋮ when seeking DUXIXENT coverage for your patients

INDICATION

DUPIXENT is indicated as an add-on maintenance treatment of adult and pediatric patients aged 6 years and older with moderate-to-severe asthma characterized by an eosinophilic phenotype or with oral corticosteroid dependent asthma.

Limitations of Use: DUXIXENT is not indicated for the relief of acute bronchospasm or status asthmaticus.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATION: Dupixent is contraindicated in patients with known hypersensitivity to dupilumab or any of its excipients.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

DUPIXENT® 
(dupilumab) Injection
200mg • 300mg

Table of contents

- 03** **Your first step: Submitting a PA request**
Guidance on how to submit the appropriate paperwork in support of your patient's access to DUPIXENT® (dupilumab)
- 05** **When a PA is denied: Navigating the appeal process**
Tips on how to compile an appeal packet if a PA is denied for DUPIXENT
- 07** **Example letters**
A list of sample letters that may provide useful information when responding to a health plan

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS

Hypersensitivity: Hypersensitivity reactions, including anaphylaxis, serum sickness or serum sickness-like reactions, angioedema, generalized urticaria, rash, erythema nodosum, and erythema multiforme have been reported. If a clinically significant hypersensitivity reaction occurs, institute appropriate therapy and discontinue DUPIXENT.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

DUPIXENT® 
(dupilumab) Injection
200mg • 300mg

Your first step: Submitting a PA request

Prior authorization is a very common requirement of health plans before approving DUPIXENT® (dupilumab). Once you have verification of an appropriate patient's pharmacy benefit provider, you should begin the PA process. You can obtain the appropriate PA form through *DUPIXENT MyWay*®, CoverMyMeds®, or your patient's insurance provider or specialty pharmacy.

» The following tips may help strengthen the case for your patient when you submit a PA on their behalf:

Document current and/or recent chart notes. This includes:

- Details of current diagnosis
- Disease severity
- Treatment history (eg, date of initial diagnosis; current and prior therapies; eosinophil, IgE, and FeNO levels and last test dates; pre-bronchodilator FEV₁ and last test date; any comorbidities; the number of severe exacerbations in the past 12 months)
- Documentation if a recommended therapy is contraindicated or inappropriate

For more tips on handling PA requirements, ask your Field Representative for a **PA checklist for asthma**



Consider including a letter that explains your patient's condition in detail. This may include a **letter of medical exception** or a **letter of medical necessity**. More information about these letters can be found on page 7.

FeNO=fractional exhaled nitric oxide; FEV₁=forced expiratory volume in 1 second; IgE= immunoglobulin E.
CoverMyMeds is a registered trademark of CoverMyMeds, LLC.

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS (cont'd)

Eosinophilic Conditions: Patients being treated for asthma may present with clinical features of eosinophilic pneumonia or eosinophilic granulomatosis with polyangiitis (EGPA). These events may be associated with the reduction of oral corticosteroid therapy. Healthcare providers should be alert to vasculitic rash, worsening pulmonary symptoms, cardiac complications, kidney injury, and/or neuropathy presenting in their patients with eosinophilia. Cases of eosinophilic pneumonia were reported in adults who participated in the asthma development program and cases of EGPA have been reported with DUPIXENT in adults who participated in the asthma development program as well as in adults with co-morbid asthma in the CRSwNP development program. Advise patients to report signs of eosinophilic pneumonia and EGPA. Consider withholding DUPIXENT if eosinophilic pneumonia or EGPA are suspected.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

DUPIXENT®
(dupilumab) Injection
200mg • 300mg

Need assistance? We are here to help

If requested on the **DUPIXENT MyWay**® Enrollment Form, the **DUPIXENT MyWay** team can provide support during the PA process, including:



Performing a **benefits investigation**



Determining **PA requirements**



Pre-populating the PA form with as much demographic information as possible



Helping to **track the PA status** with the patient's health plan and communicating with you and your patient about the status

For additional information or if you have questions, contact your Field Representative or call **DUPIXENT MyWay** at **1-844-DUPIXEN(T)** (1-844-387-4936) Option 1, Monday–Friday, 8 AM–9 PM Eastern time.

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS (cont'd)

Acute Symptoms of Asthma or Acute Deteriorating Disease: Do not use DAPIXENT to treat acute symptoms or acute exacerbations of asthma, acute bronchospasm, or status asthmaticus. Patients should seek medical advice if their asthma remains uncontrolled or worsens after initiation of DAPIXENT.

Risk Associated with Abrupt Reduction of Corticosteroid Dosage: Do not discontinue systemic, topical, or inhaled corticosteroids abruptly upon initiation of DAPIXENT. Reductions in corticosteroid dose, if appropriate, should be gradual and performed under the direct supervision of a healthcare provider. Reduction in corticosteroid dose may be associated with systemic withdrawal symptoms and/or unmask conditions previously suppressed by systemic corticosteroid therapy.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

DUPIXENT®
(dupilumab) Injection
200mg • 300mg

When PA is denied: Navigating the appeal process

If your patient's PA is denied, you can appeal the decision. A successful appeal may include the following checklist of information, compiled in an appeal packet.

- ☐ A letter of appeal signed by the treating physician and patient or caregiver, if required. Appeal letters can be customized depending on the reason a PA has been denied. See page 7 for a list of example letters that you can download and reference to appeal a PA denial
- ☐ The appeal form recommended by the health plan
- ☐ In addition to the letter of appeal and appeal form, consider including current and/or recent chart notes from the patient's treating physician to make the appeal as thorough as possible, including:
 - ☐ Date of initial diagnosis
 - ☐ Eosinophil levels (for eosinophilic phenotype) and last test date
 - ☐ FeNO levels (if available) and last test date
 - ☐ IgE levels (if available) and last test date
 - ☐ Pre-bronchodilator forced expiratory volume in 1 second (FEV₁) and last test date
 - ☐ Number of severe exacerbations in the past 12 months
 - ☐ Level of asthma control
 - ☐ Oral corticosteroid use
 - ☐ Maximally tolerated inhaled corticosteroid dose
 - ☐ Response to all prior therapies
 - ☐ Any relevant comorbidities or contraindications
 - ☐ If appropriate, earlier treatment history from previous physicians, provided by the patient
- ☐ *International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM)* code matching your patient's diagnosis
- ☐ Reasons why the patient's recent symptoms, severity of condition, and impact of disease warrant treatment with DUPIXENT® (dupilumab)
- ☐ Any clinical studies or peer-reviewed articles documenting the medical effectiveness of DUPIXENT
- ☐ DUPIXENT full Prescribing Information, available at www.DUPIXENThcp.com
- ☐ A personal narrative from the patient that describes the impact of their condition

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS (cont'd)

Psoriasis: Cases of new-onset psoriasis have been reported with the use of DUPIXENT for the treatment of asthma, including in patients without a family history of psoriasis. In postmarketing reports, these cases resulted in partial or complete resolution of psoriasis with discontinuation of dupilumab, with or without use of supplemental treatment for psoriasis (topical or systemic). Those who continued dupilumab received supplemental treatment for psoriasis to improve associated symptoms. Advise patients to report new-onset psoriasis symptoms. If symptoms persist or worsen, consider dermatologic evaluation and/or discontinuation of DUPIXENT.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

DUPIXENT
(dupilumab) Injection
200mg • 300mg

When PA is denied: Navigating the appeal process (cont'd)

Key points to consider when filing an appeal on behalf of your patient:

- Adhere to the timelines and use the forms noted in the health plan's letter of denial
- Depending on the health plan, your patient's signature may be required on the appeal letter (if patient is a minor, a guardian's signature is required)
- The appeal packet should be submitted by your office or your patient
- Two levels of internal review may be required before the health plan will notify you of your patient's eligibility for an external appeal. If this occurs, the reviewer will be an independent party, typically board certified in the specialty. Their decision will be binding on the health plan
 - All documentation from previous reviews should be submitted in subsequent appeals

Remember, successful appeals may take more than 1 attempt.¹ Patients can also advocate for an appeal on their own behalf, and HCPs may request a peer-to-peer review with a medical reviewer at a health plan.

DUPIXENT MyWay® can help educate your office about the appropriate actions needed to appeal a coverage denial. DUXIPENT MyWay Appeal Specialists can help provide support throughout the appeal process.

For additional information or if you have questions, contact your Field Representative or call **DUPIXENT MyWay** at **1-844-DUXIPENT(T)** (1-844-387-4936) Option 1, Monday–Friday, 8 AM–9 PM Eastern time.

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS (cont'd)

Arthralgia and Psoriatic Arthritis: Arthralgia has been reported with use of DUXIPENT with some patients reporting gait disturbances or decreased mobility associated with joint symptoms; some cases resulted in hospitalization. Cases of new-onset psoriatic arthritis requiring systemic treatment have been reported with the use of DUXIPENT. Advise patients to report new onset or worsening joint symptoms. If symptoms persist or worsen, consider rheumatological evaluation and/or discontinuation of DUXIPENT.

Parasitic (Helminth) Infections: It is unknown if DUXIPENT will influence the immune response against helminth infections. Treat patients with pre-existing helminth infections before initiating therapy with DUXIPENT. If patients become infected while receiving treatment with DUXIPENT and do not respond to anti-helminth treatment, discontinue treatment with DUXIPENT until the infection resolves. Helminth infections (5 cases of enterobiasis and 1 case of ascariasis) were reported in pediatric patients 6 to 11 years old in the pediatric asthma development program.

Vaccinations: Consider completing all age-appropriate vaccinations as recommended by current immunization guidelines prior to initiating DUXIPENT. Avoid use of live vaccines during treatment with DUXIPENT.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

DUPIXENT®
(dupilumab) Injection
200mg • 300mg

Example letters

The following example letters are templates for the information that may be required when responding to a PA or appeal request for DUPIXENT® (dupilumab) from a patient's health plan. Use of the information within these letters does not guarantee that the health plan will provide reimbursement for DUPIXENT and is not intended to be a substitute for or to influence the independent medical judgment of the physician.



Sample letter of medical necessity

Consider including this letter: To emphasize that DUPIXENT was prescribed because it is necessary for the patient's health and will result in better outcomes. This letter can be accompanied with a PA submission or in addition to your appeal letter, if needed

Who should sign this letter: HCP only



Sample medical exception letter

Consider including this letter: If coverage for DUPIXENT is denied because of the health plan's policy or if DUPIXENT is subject to a national drug code block. This letter can be accompanied with a PA submission or in addition to your appeal letter, if needed

Who should sign this letter: Both the patient or caregiver and HCP



Denial due to severity

Consider including this letter: If coverage for DUPIXENT is denied because your patient's condition did not meet the plan's severity criteria for treatment with DUPIXENT

Who should sign this letter: Both the patient or caregiver and HCP



Sample appeal letter for denial due to nonformulary status

Consider including this letter: If coverage is denied because DUPIXENT is not on the health plan's formulary or not covered for any other reason

Who should sign this letter: Both the patient or caregiver and HCP

[Click here to download these example letters](#)

IMPORTANT SAFETY INFORMATION (cont'd)

ADVERSE REACTIONS: The most common adverse reactions (incidence $\geq 1\%$) in patients with asthma are injection site reactions, oropharyngeal pain, and eosinophilia.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

DUPIXENT
(dupilumab) Injection
200mg • 300mg



For more information, contact your
Field Representative or call *DUPIXENT MyWay*® at
1-844-DUPIXEN(T) (1-844-387-4936) Option 1,
Monday–Friday, 8 AM–9 PM Eastern time

Reference: 1. United States Government Accountability Office. Report to the Secretary of Health and Human Services and the Secretary of Labor. Private health insurance: data on application and coverage denials. March 2011. Accessed October 25, 2024. <https://www.gao.gov/assets/320/316699.pdf>

IMPORTANT SAFETY INFORMATION (cont'd)

USE IN SPECIFIC POPULATIONS

- **Pregnancy:** A pregnancy exposure registry monitors pregnancy outcomes in women exposed to DUPIXENT during pregnancy. Available data from case reports and case series with DUPIXENT use in pregnant women have not identified a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. Human IgG antibodies are known to cross the placental barrier; therefore, DUPIXENT may be transmitted from the mother to the developing fetus.
- **Lactation:** There are no data on the presence of DUPIXENT in human milk, the effects on the breastfed infant, or the effects on milk production. Maternal IgG is known to be present in human milk. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for DUPIXENT and any potential adverse effects on the breastfed child from DUPIXENT or from the underlying maternal condition.

Please see accompanying full [Prescribing Information](#).